

University of Groningen

Discovery, characterization, and kinetic analysis of an alditol oxidase from streptomyces coelicolor

Heuts, Dominic P. H. M.; van Hellemond, Erik W.; Janssen, Dick B.; Fraaije, Marco W.

Published in:
The Journal of Biological Chemistry

DOI:
[10.1074/jbc.M610849200](https://doi.org/10.1074/jbc.M610849200)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Heuts, D. P. H. M., van Hellemond, E. W., Janssen, D. B., & Fraaije, M. W. (2007). Discovery, characterization, and kinetic analysis of an alditol oxidase from streptomyces coelicolor. *The Journal of Biological Chemistry*, 282(28), 20283-20291. <https://doi.org/10.1074/jbc.M610849200>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

APPENDIX

DISCOVERY, CHARACTERIZATION AND KINETIC ANALYSIS OF AN ALDITOL OXIDASE FROM *STREPTOMYCES COELICOLOR*

Dominic P.H.M Heuts, Erik W. van Hellemond, Dick B. Janssen, Marco W. Fraaije
From the Laboratory of Biochemistry, Groningen Biomolecular Sciences and Biotechnology Institute,
University of Groningen, The Netherlands


Running title: Alditol oxidase from *S. coelicolor*

Correspondence to: Dr. M.W. Fraaije, Laboratory of Biochemistry, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands. Tel: + 31 50 3634345; Fax: + 31 50 36 34165; E-Mail: m.w.fraaije@rug.nl

Steady-state kinetics

We used the determinant method for obtaining the steady-state rate equations of the proposed ping-pong and ternary complex mechanisms (1). From these rate equations the steady-state kinetic parameters were calculated. Next to this, we also show the measured and simulated Lineweaver-Burk plots of steady-state kinetics of AldO at varying oxygen and xylitol concentrations. The plots and the calculated steady-state kinetic parameters suggest that AldO follows a ternary complex mechanism.

Scheme 1. Proposed kinetic scheme (I = ping-pong mechanism, II = ternary complex mechanism).



Ping-pong mechanism

To solve the steady-state rate equation for the proposed ping-pong mechanism, the steady-state concentrations of all enzyme species need to be solved. This is accomplished by using the determinant method (1). Because of the relatively fast equilibrium between E_{ox} and $E_{ox} \sim S$ these species are lumped yielding: $d(E_{ox} + E_{ox} \sim S)/dt = -[E_{ox} \sim S] \cdot k_{red} + [E_{red}] \cdot k_{ox, 1} \cdot O_2$ where $[E_{ox} \sim S] = ([E_{ox}] + [E_{ox} \sim S]) \cdot S / (K_d + S)$. Substitution of $[E_{ox} \sim S]$ by the latter equation then yields:

$$d(E_{ox} + E_{ox} \sim S)/dt = -([E_{ox}] + [E_{ox} \sim S]) \cdot S / (K_d + S) \cdot k_{red} + [E_{red}] \cdot k_{ox, 1} \cdot O_2.$$

	$E_{ox} + E_{ox} \sim S$	$E_{red} \sim P'$	$E_{red} \sim P$	E_{red}
$d(E_{ox} + E_{ox} \sim S)/dt$	$-k_{red} * \frac{S}{K_d + S}$	0	0	$k_{ox,1} * O_2$
$d(E_{red} \sim P')/dt$	$k_{red} * \frac{S}{K_d + S}$	$-k_3$	k_{-3}	0
$d(E_{red} \sim P)/dt$	0	k_3	$-k_4 - k_{-3}$	0
$d(E_{red})/dt$	0	0	k_4	$-k_{ox,1} * O_2$

From this matrix the concentration for ($E_{ox} + E_{ox} \sim S$) is obtained by deleting the first row and first column and then calculating the determinant of the remaining smaller matrix. In a similar way the concentrations of the remaining enzyme species can be obtained. These values can then be used to write down the steady-state rate equation in the form of a Michaelis-Menten equation:

$$V = \frac{k_{ox,1} O_2 [E_{red}]}{[E_{ox} + E_{ox} \sim S] + [E_{red} \sim P'] + [E_{red} \sim P] + [E_{red}]}$$

$$V = \frac{k_3 k_4 k_{ox,1} k_{red} O_2 S}{(k_4 + k_{-3}) k_{ox,1} k_{red} O_2 S + k_3 (k_{ox,1} k_{red} O_2 S + k_4 (K_d k_{ox,1} O_2 + k_{red} S + k_{ox,1} O_2 S))}$$

From this rate equation the kinetic parameters K_M and k_{cat} can be derived:

$$K_M = \frac{k_3 k_4 K_d k_{ox,1} O_2}{(k_4 + k_{-3}) k_{ox,1} k_{red} O_2 + k_3 (k_{ox,1} k_{red} O_2 + k_4 (k_{red} + k_{ox,1} O_2))}$$

$$k_{cat} = \frac{k_3 k_4 k_{ox,1} k_{red} O_2}{(k_4 + k_{-3}) k_{ox,1} k_{red} O_2 + k_3 (k_{ox,1} k_{red} O_2 + k_4 (k_{red} + k_{ox,1} O_2))}$$

$$\frac{k_{cat}}{K_M} = \frac{k_{red}}{K_d}$$

Ternary complex mechanism

The steady-state rate equation for the proposed ternary complex mechanism can be derived in a similar way. The result is:

$$V = \frac{k_3 k_7 k_{ox,2} k_{red} O_2 S}{(k_7 k_{red} (k_{-3} + k_{ox,2} O_2) S + k_3 (k_{ox,2} k_{red} O_2 S + k_7 (K_d k_{ox,2} O_2 + k_{red} S + k_{ox,2} O_2 S)))}$$

From this rate equation the following equations for K_M and k_{cat} were derived:


$$K_M = \frac{k_3 k_7 K_d k_{ox,2} O_2}{k_7 k_{red} (k_{-3} + k_{ox,2} O_2) + k_3 (k_{red} k_{ox,2} O_2 + k_7 (k_{red} + k_{ox,2} O_2))}$$

$$k_{cat} = \frac{k_3 k_7 k_{red} k_{ox,2} O_2}{k_7 k_{red} (k_{-3} + k_{ox,2} O_2) + k_3 (k_{red} k_{ox,2} O_2 + k_7 (k_{red} + k_{ox,2} O_2))}$$

$$\frac{k_{cat}}{K_M} = \frac{k_{red}}{K_d}$$

Steady-state kinetics of AldO at varying oxygen concentrations

Steady-state kinetic parameters of AldO were determined at 0.25 and 1.25 mM O_2 (k_{cat} is respectively 13 s^{-1} and 20 s^{-1} , K_M is respectively 0.32 mM and 0.49 mM). The set of Lineweaver-Burke plots obtained in this way gives information on the kinetic mechanism that is operative. A set of parallel lines suggests that a ping-pong mechanism is employed by the enzyme, however it is known that in certain cases a ternary complex mechanism yields the same result (2,3). For AldO a set of parallel lines was obtained and correlated to a ternary complex mechanism by simulating the Lineweaver-Burk plot according to the abovementioned equations.



Tested compounds for substrate profiling

D-ribose, L-arginine, diglycerol, L-arabinose, L-alanine, cholesterol, D-xylose, D-alanine, D-lyxose, L-asparagine, D-glucose, L-aspartate, ethyleneglycol, D-mannose, DL-aspartate, diethyleneglycol, D-galactose, L-cysteine, hexaethyleneglycol, L-histidine, PEG, L-proline, D-fructose, L-threonine, 3-buten-1-ol, 3-buten-2-ol, glycerol, sarcosine, cis-2-Butene-1,4-diol, meso-erythritol, DL-homoserine, L-threitol, L-ornithine, ribitol, DL-norvaline, 2-aminoethanol, D-arabitol, 1-amino-2-propanol, xylitol, 2-amino-1-propanol, D-sorbitol, methanol, 4-amino-1-butanol, D-mannitol, ethanol, galactitol, 1-propanol, 2-propanol, 1-butanol, 1,4-diaminobutane, L-rhamnose, 2-butanol, L-fucose, 1-pentanol, 2-pentanol, benzyl alcohol, 1-octanol, 2-phenylethanol, L-gulonolactone, isoamylalcohol, α -methylbenzyl alcohol, α -methylbenzyl amine, 1-phenyl-1,2-ethanediol, maltose, 1,2-propanediol, 2-amino-1-phenylethanol, lactose, 1,3-propanediol, 2-amino-2-phenylethanol, sucrose, 1,2-butanediol, 2-amino-2-phenylethanol, 1,3-butanediol benzyl methylether, 1,4-butanediol, cinnamyl alcohol, D-melizitol, meso-2,3-butanediol, 2-methoxy-4-methylphenol, N-acetylglucosamine, butanediol, vanillyl alcohol, butanediol, 1,2-pentanediol, 1,2-hexanediol, cis-1,2-cyclohexanediol.

REFERENCES

1. Huang, C. Y.(1983) in *Contemporary enzyme kinetics and mechanism*, Academic Press, New York, 54-84
2. Porter, D. J., Voet, J. G., and Bright, H. J. (1977) *J. Biol. Chem.* **252**, 4464-4473
3. Fraaije, M. W. and Van Berkel, W. J. (1997) *J. Biol. Chem.* **272**, 18111-18116